

phone: (206) 543-8637; fax: (206) 616-5927 e-mail: naccmail@u.washington.edu website: www.alz.washington.edu

NACC Uniform Data Set (UDS) - Initial Visit Packet

Form B3: Evaluation Form – Unified Parkinson's Disease Rating Scale (UPDRS¹) – Motor Exam

Center:		ADC Subject ID:			Form Date://		
For additional clarific			to be completed by the clinician or other trained heal cation and examples, see UDS Coding Guidebook fo eck only <u>one</u> box per question.				
	[Optional] end form h		clinician completes the UPDRS examination	on and	determines all items are normal, check this box and		
1.	Speech	□ 1	Normal. Slight loss of expression, diction and/or volume. Monotone, slurred but understandable; moderately impaired.	□ 4	Marked impairment, difficult to understand. Unintelligible. Untestable (specify reason):		
2.	Facial expression	□ 1	Normal. Minimal hypomimia, could be normal "poker face". Slight but definitely abnormal diminution of facial expression.	□ 4	Moderate hypomimia; lips parted some of the time. Masked or fixed facies with severe or complete loss of facial expression; lips parted ¼ inches or more. Untestable (specify reason):		
3.	. Tremor at rest						
3a.	Face, lips, chin	□ 1	Absent. Slight and infrequently present. Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present.	□ 4	Moderate in amplitude and present most of the time. Marked in amplitude and present most of the time. Untestable (<i>specify reason</i>):		
3b.	Right hand	□ 1	Absent. Slight and infrequently present. Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present.	□ 4	Moderate in amplitude and present most of the time. Marked in amplitude and present most of the time. Untestable (<i>specify reason</i>):		

(version 2.0, February 2008)

Fahn S, Elton RL, UPDRS Development Committee. The Unified Parkinson's Disease Rating Scale. In Fahn S, Marsden CD, Calne DB, Goldstein M, eds. Recent developments in Parkinson's disease, Vol. 2. Florham Park, NJ: Macmillan Healthcare Information, 1987:153-163, 293-304. Reproduced by permission of the author.

C C 11 C	er:		ADC Subject ID:		Form Date://	
For a	NOTE: This form is to be completed by the clinician or other trained health professional. ADC Visit #:					
3c.	Left hand		Absent.	□ 3	Moderate in amplitude and present most of the time.	
	nana		Slight and infrequently present.	$\Box 4$	Marked in amplitude and present most of the time.	
		□ 2	Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present.		Untestable (specify reason):	
3d.	Right	□ 0	Absent.	□ 3	Moderate in amplitude and present most of the	
	foot	\Box 1	Slight and infrequently present.		time.	
		\square 2	Mild in amplitude and persistent; or		Marked in amplitude and present most of the time.	
			moderate in amplitude, but only intermittently present.	□ 8	Untestable (specify reason):	
3e.	Left		Absent.	□ 3	Moderate in amplitude and present most of the	
	foot	\Box 1	Slight and infrequently present.		time.	
		\square 2	Mild in amplitude and persistent; or		Marked in amplitude and present most of the time.	
			moderate in amplitude, but only intermittently present.	□ 8	Untestable (specify reason):	
4.	Action or p	ostura	al tremor of hands			
	Action or p		Absent.	□ 3	Moderate in amplitude with posture holding as well	
				□ 3	Moderate in amplitude with posture holding as well as action.	
	Right	□ 0 □ 1	Absent. Slight; present with action. Moderate in amplitude, present with	□ 4	Marked in amplitude; interferes with feeding.	
	Right	□ 0 □ 1	Absent. Slight; present with action.	□ 4	as action.	
4a.	Right hand	□ 0 □ 1 □ 2	Absent. Slight; present with action. Moderate in amplitude, present with	□ 4 □ 8	as action. Marked in amplitude; interferes with feeding. Untestable (<i>specify reason</i>): Moderate in amplitude with posture holding as well	
4a.	Right hand	□ 0 □ 1 □ 2	Absent. Slight; present with action. Moderate in amplitude, present with action.	□ 4 □ 8	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action.	
4a.	Right hand	□ 0 □ 1 □ 2 □ 0 □ 1	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action. Moderate in amplitude, present with	□ 4 □ 8 □ 3 □ 4	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action. Marked in amplitude; interferes with feeding.	
4a.	Right hand	□ 0 □ 1 □ 2 □ 0 □ 1	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action.	□ 4 □ 8 □ 3 □ 4	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action.	
4a. 4b.	Right hand Left hand	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action. Moderate in amplitude, present with action.	□ 4 □ 8 □ 3 □ 4 □ 8	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action. Marked in amplitude; interferes with feeding. Untestable (specify reason):	
4a. 4b.	Right hand Left hand	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action. Moderate in amplitude, present with action.	□ 4 □ 8 □ 3 □ 4 □ 8	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action. Marked in amplitude; interferes with feeding.	
4a. 4b.	Right hand Left hand Rigidity (ju	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2 □ 1 □ 2	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action. Moderate in amplitude, present with action.	□ 4 □ 8 □ 3 □ 4 □ 8	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action. Marked in amplitude; interferes with feeding. Untestable (specify reason):	
4a. 4b.	Right hand Left hand Rigidity (juignored)	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2 □ 1 □ 2	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action. Moderate in amplitude, present with action. on passive movement of major joints with Absent. Slight or detectable only when activated	□ 4 □ 8 □ 3 □ 4 □ 8 patien	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): t relaxed in sitting position; cogwheeling to be	
4a. 4b.	Right hand Left hand Rigidity (juignored)	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2 □ 1 □ 2 □ 1 □ 1 □ 1	Absent. Slight; present with action. Moderate in amplitude, present with action. Absent. Slight; present with action. Moderate in amplitude, present with action. on passive movement of major joints with Absent.	□ 4 □ 8 □ 3 □ 4 □ 8 patien	as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): Moderate in amplitude with posture holding as well as action. Marked in amplitude; interferes with feeding. Untestable (specify reason): t relaxed in sitting position; cogwheeling to be Marked, but full range of motion easily achieved.	

Cent	ter:		ADC Subject ID:		Form Date://	
For	NOTE: This form is to be completed by the clinician or other trained health professional. ADC Visit #:					
5b.	Right	$\Box 0$	Absent.	□ 3	Marked, but full range of motion easily achieved.	
	upper	\Box 1	Slight or detectable only when activated by mirror or other movements.	□ 4	Severe; range of motion achieved with difficulty.	
	extremity			□ 8	Untestable (specify reason):	
		□ 2	Mild to moderate.			
5c.	Left	\square 0	Absent.	□ 3	Marked, but full range of motion easily achieved.	
	upper extremity		by mirror or other movements.	□ 4	Severe; range of motion achieved with difficulty.	
	extremity			□ 8	Untestable (specify reason):	
		$\frac{\sqcup 2}{}$	Mild to moderate.			
5d.	Right	\square 0	Absent.	□ 3	Marked, but full range of motion easily achieved.	
	lower extremity	\Box 1	Slight or detectable only when activated	□ 4	Severe; range of motion achieved with difficulty.	
	CAUCHILI		by mirror or other movements.	□ 8	Untestable (specify reason):	
			Mild to moderate.			
5e.	Left	$\Box 0$	Absent.		Marked, but full range of motion easily achieved.	
	lower extremity	\Box 1	Slight or detectable only when activated		Severe; range of motion achieved with difficulty.	
		\Box 2	by mirror or other movements. Mild to moderate.	□ 8	Untestable (specify reason):	
			Willa to illouerate.			
6.	Finger taps	(patie	ent taps thumb with index finger in rapid su	ccessi	on)	
	Right	_	ent taps thumb with index finger in rapid su Normal.		Severely impaired; frequent hesitation in initiating	
			Normal. Mild slowing and/or reduction in	□ 3	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement.	
	Right	□ 0 □ 1	Normal. Mild slowing and/or reduction in amplitude.	□ 3 □ 4	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task.	
	Right	□ 0 □ 1	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early	□ 3 □ 4	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement.	
	Right	□ 0 □ 1	Normal. Mild slowing and/or reduction in amplitude.	□ 3 □ 4	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task.	
ба.	Right	□ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in	□ 3 □ 4 □ 8	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task.	
ба.	Right hand	□ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement.	□ 3 □ 4 □ 8	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>):	
ба.	Right hand		Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal.	□ 3 □ 4 □ 8	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>): Severely impaired; frequent hesitation in initiating	
ба.	Right hand	□ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early	□ 3 □ 4 □ 8 □ 3 □ 4	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement.	
ба.	Right hand	□ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude.	□ 3 □ 4 □ 8 □ 3 □ 4	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task.	
ба.	Right hand	□ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in	□ 3 □ 4 □ 8 □ 3 □ 4	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task.	
6a.	Right hand Left hand		Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in	□ 3 □ 4 □ 8 □ 3 □ 4 □ 8	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (<i>specify reason</i>):	
6a. 6b.	Right hand Left hand Hand move	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2 □ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement.	□ 3 □ 4 □ 8 □ 3 □ 4 □ 8 □ 4 □ 8	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (specify reason): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (specify reason): ion) Severely impaired; frequent hesitation in initiating	
6a. 6b.	Right hand Left hand Hand move	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2 □ 0 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. s (patient opens and closes hands in rapid so Normal. Mild slowing and/or reduction in	□ 3 □ 4 □ 8 □ 3 □ 4 □ 8 □ 3	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (specify reason): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (specify reason): ion) Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement.	
6a. 6b.	Right hand Left hand Hand move	□ 0 □ 1 □ 2 □ 0 □ 1 □ 2 □ 0 □ 1 □ 1 □ 2	Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. Normal. Mild slowing and/or reduction in amplitude. Moderately impaired; definite and early fatiguing; may have occasional arrests in movement. s (patient opens and closes hands in rapid survey).	□ 3 □ 4 □ 8 □ 3 □ 4 □ 8 □ 3 □ 4 □ 8	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (specify reason): Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement. Can barely perform the task. Untestable (specify reason): ion) Severely impaired; frequent hesitation in initiating	

Center:		ADC Subject ID:		Form Date://		
NOTE: This form is to be completed by the clinician or other trained health professional. ADC Visit #: For additional clarification and examples, see UDS Coding Guidebook for Initial Visit Packet, Form B3.Check only <u>one</u> box per question.						
7b. Left hand		Normal. Mild slowing and/or reduction in	□ 3	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement.		
		amplitude.	□ 4	Can barely perform the task.		
	□ 2	Moderately impaired; definite and early fatiguing; may have occasional arrests in movement.	□ 8	Untestable (specify reason):		
	8. Rapid alternating movements of hands (pronation-supination movements of hands, vertically and horizontally, with as large an amplitude as possible, both hands simultaneously)					
8a. Right \square 0 Normal. hand \square 1 Mild slowing and/or reduce		Normal. Mild slowing and/or reduction in	□ 3	Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement.		
		amplitude.	□ 4	Can barely perform the task.		
	□ 2	Moderately impaired; definite and early fatiguing; may have occasional arrests in movement.	□ 8	Untestable (specify reason):		
8b. Left □ 0 Normal.		□ 3	Severely impaired; frequent hesitation in initiating			
hand	□ 1	Mild slowing and/or reduction in		movements or arrests in ongoing movement.		
		amplitude.		Can barely perform the task.		
	□ 2	Moderately impaired; definite and early fatiguing; may have occasional arrests in movement.	□ 8	Untestable (specify reason):		
9. Leg agility (patient taps heel on the ground in rapid succession, picking up entire leg; amplitude should be at least 3 inches)						
9a. Right leg	$\square 0$	Normal.	□ 3	Severely impaired; frequent hesitation in initiating		
	\Box 1	Mild slowing and/or reduction in		movements or arrests in ongoing movement.		
		amplitude.		Can barely perform the task.		
	□ 2	Moderately impaired; definite and early fatiguing; may have occasional arrests in movement.	⊔ 8	Untestable (specify reason):		
9b. Left leg	□ 0	Normal.	□ 3	Severely impaired; frequent hesitation in initiating		
	□ 1	Mild slowing and/or reduction in		movements or arrests in ongoing movement.		
	\Box 2	amplitude. Moderately impaired; definite and early		Can barely perform the task. Untestable (<i>specify reason</i>):		
	⊔ ∠	fatiguing; may have occasional arrests in movement.	⊔ 0			

Center:	ADC Subject ID:	Form Date://
	eted by the clinician or other trained health examples, see UDS Coding Guidebook for l e box per question.	
10. Arising from chair (patient attempts to rise from a straight-backed chair, with arms folded across chest)	□ 1 Slow; or may need more than one attempt.□ 2 Pushes self up from arms of seat.	 □ 3 Tends to fall back and may have to try more than one time, but can get up without help. □ 4 Unable to arise without help. □ 8 Untestable (specify reason): □
11. Posture	 □ 1 Not quite erect, slightly stooped posture; could be normal for older person. □ 2 Moderately stooped posture, 	 □ 3 Severely stooped posture with kyphosis; can be moderately leaning to one side. □ 4 Marked flexion with extreme abnormality of posture. □ 8 Untestable (specify reason):
12. Gait	☐ 1 Walks slowly; may shuffle with short steps, but no festination (hastening steps) or propulsion.	 □ 3 Severe disturbance of gait requiring assistance. □ 4 Cannot walk at all, even with assistance. □ 8 Untestable (specify reason):
13. Posture stability (response to sudden, strong posterior displacement produced by pull on shoulders while patient erect with eyes open and feet slightly apart; patient is prepared)	 □ 1 Retropulsion, but recovers unaided. □ 2 Absence of postural response; would fall if not caught by 	 □ 3 Very unstable, tends to lose balance spontaneously. □ 4 Unable to stand without assistance. □ 8 Untestable (specify reason):
14. Body bradykinesia and hypokinesia (combining slowness, hesitancy, decreased arm swing, small amplitude, and poverty of movement in general)	☐ 1 Minimal slowness, giving movement a deliberate character; could be normal for some persons;	 □ 3 Moderate slowness, poverty or small amplitude of movement. □ 4 Marked slowness, poverty or small amplitude of movement. □ 8 Untestable (<i>specify reason</i>):